

A Minimal Groebner Basis for the Defining Ideals of Certain Affine Monomial Curves

Ibrahim Al-Ayyoub

February 1, 2008

Abstract

Let K be a field and let $m_0, \dots, m_n$ be an almost arithmetic sequence of positive integers. Let C be a monomial curve in the affine $(n+1)$ -space, defined parametrically by $x_0 = t^{m_0}, \dots, x_n = t^{m_n}$. In this article we produce Groebner bases for the defining ideal of C , correcting previous work of Sengupta, (2003).

Introduction

Let $n \geq 2$, K a field and let $x_0, \dots, x_n, t$ be indeterminates. Let $m_0, \dots, m_n$ be an almost arithmetic sequence of positive integers, that is, some $n-1$ of these form an arithmetic sequence, and assume $\gcd(m_0, \dots, m_n) = 1$. Let P be the kernel of the K -algebra homomorphism $\eta : K[x_0, \dots, x_n] \rightarrow K[t]$, defined by $\eta(x_i) = t^{m_i}$. A set of generators for the ideal P was explicitly constructed in Patil and Singh (1990). We call these generators the “*Patil-Singh generators*”. Out of this generating set, Patil (1993) constructed a minimal set Ω for the ideal P . We call the elements of Ω “*Patil generators*”. Sengupta (2003) proved that Ω forms a Groebner basis for the relation ideal P with respect to the grevlex monomial order, however, Sengupta’s proof is not complete, as in fact Ω is not a Groebner basis in all cases, see Remark 1.6 and Remark 1.7. The goal of this article is to produce a minimal Groebner basis for P . Remark 1.6 was a motive to modify Patil generators before computing Groebner basis. In Section 2 the set Ω is described more explicitly and it is modified. Then in Section 3 we state and prove a general result that we use to produce a Groebner basis. This makes the proof of the main result of this thesis much shorter and simpler than the work given in Sengupta (2003).

1 Patil Generators

We shall use the notation and the terminology from Patil and Singh (1990) and Patil (1993) with a slight difference in naming some variables and constants. Let $n \geq 2$ be an integer and let $p = n - 1$. Let $m_0, \dots, m_p$ be an arithmetic sequence of positive integers with $0 < m_0 < \dots < m_p$, let m_n be arbitrary, and $\gcd(m_0, \dots, m_n) = 1$. Let Γ denote the numerical semigroup that is generated by $m_0, \dots, m_n$ i.e. $\Gamma = \sum_{i=0}^n \mathbb{N}_0 m_i$. We assume throughout that Γ is minimally generated by $m_0, \dots, m_n$.

Put $\Gamma' = \sum_{i=0}^p \mathbb{N}_0 m_i$. Thus $\Gamma = \Gamma' + \mathbb{N}_0 m_n$.

Notation 1.1 For $a, b \in \mathbb{Z}$ let $[a, b] = \{t \in \mathbb{Z} \mid a \leq t \leq b\}$. For $t \geq 0$, let $q_t \in \mathbb{Z}$, $r_t \in [1, p]$ and $g_t \in \Gamma'$ be defined by $t = q_t p + r_t$ and $g_t = q_t m_p + m_{r_t}$.

Let $S = \{\gamma \in \Gamma \mid \gamma - m_0 \notin \Gamma\}$. As all large integers are in Γ , S is a finite set of non-negative integers. The following gives an explicit description of S .

Lemma 1.2 (Lemma 3.1, 3.2 in Patil-Singh (1990)) Let $u = \min\{t \geq 0 \mid g_t \notin S\}$ and $v = \min\{b \geq 1 \mid b m_n \in \Gamma'\}$.

(a) There exist unique integers $w \in [0, v-1]$, $z \in [0, u-1]$, $\lambda \geq 1$, $\mu \geq 0$, and $\nu \geq 2$ such that

$$(i) \quad g_u = \lambda m_0 + w m_n;$$

$$(ii) \quad v m_n = \mu m_0 + g_z;$$

$$(iii) \quad g_{u-z} + (v-w)m_n = \nu m_0, \text{ where } \nu = \begin{cases} \lambda + \mu + 1, & \text{if } r_{u-z} < r_u; \\ \lambda + \mu, & \text{if } r_{u-z} \geq r_u. \end{cases}$$

(b) Let $V = [0, u-1] \times [0, v-1]$ and $W = [u-z, u-1] \times [v-w, v-1]$. Then every element of Γ can be expressed uniquely in the form $a m_0 + g_s + b m_n$ with $a \in \mathbb{N}_0$ and $(s, b) \in V - W$.

Proof. The proof can be found in Patil and Singh (1990). We would like to recall their proof for part (b) which constitutes of a major point in our proof of the main theorem of this article. The existence of $a \in \mathbb{N}_0$ and $(s, b) \in V - W$ such that every element of Γ can be expressed in the form $a m_0 + g_s + b m_n$ is a consequence of the main proposition of this article, Proposition 3.6. Here we prove uniqueness. Let $\gamma \in \Gamma$ such that $\gamma = a m_0 + g_s + b m_n$ and also $\gamma = a' m_0 + g_t + c m_n$ with $(s, b), (t, c) \in V - W$. This implies $g_s + b m_n \equiv g_t + c m_n \pmod{m_0}$. Assume $b \geq c$, then $0 \leq (b-c)m_n \equiv g_{s-t} \pmod{m_0}$, hence we also must assume that $s \leq t$. We consider two cases: let $e \geq 0$. Case $(b-c)m_n = g_{s-t} + e m_0$ is a contradiction since $b-c < v$. Case $(b-c)m_n + e m_0 = g_{s-t}$. If $e > 0$ then $g_{s-t} - m_0 = (b-c)m_n + (e-1)m_0 \in \Gamma$, but this is a contradiction since $s-t < u$. Thus, $e = 0$. Therefore, $(b-c)m_n = g_{s-t}$. But $b-c < v$ and $s-t < u$ by assumption. Hence, $b = c$ and $s = t$. ■

Notation 1.3 Let $q = q_u$, $r = r_u$, $q' = q_{u-z}$, $r' = r_{u-z}$. From now on, the symbols q , q' , r , r' , u , v , w , z , λ , μ , ν , V and W will have the meaning assigned to them by this notation and the lemma above.

Remark 1.4 Note that for $1 \leq i \leq p$ we have $g_i - m_0 = m_i - m_0$. Then by the minimality assumption on the generators of Γ it follows that $u > p$, hence $q > 0$.

We recall the construction and the result given in Patil (1993): recall $p = n - 1$ and let

$$\begin{aligned} \xi_{i,j} &= \begin{cases} x_i x_j - x_0 x_{i+j}, & \text{if } i+j \leq p; \\ x_i x_j - x_{i+j-p} x_p, & \text{if } i+j > p, \end{cases} \\ \varphi_i &= x_{r+i} x_p^q - x_0^{\lambda-1} x_i x_n^w, \\ \psi_j &= x_{r'+j} x_p^{q'} x_n^{v-w} - x_0^{\nu-1} x_j, \\ \theta &= \begin{cases} x_n^v - x_0^\mu x_{r-r'} x_p^{q-q'}, & \text{if } r' < r; \\ x_n^v - x_0^\mu x_{p+r-r'} x_p^{q-q'-1}, & \text{if } r' \geq r, \end{cases} \end{aligned}$$

Let

$$\begin{aligned} I &= \begin{cases} [0, p-r], & \text{if } \mu \neq 0 \text{ or } W = \phi; \\ [\max(r_z - r + 1, 0), p-r], & \text{if } \mu = 0 \text{ and } W \neq \phi, \end{cases} \\ J &= \begin{cases} \phi, & \text{if } W = \phi; \\ [0, \min(z-1, p-r')], & \text{if } W \neq \phi. \end{cases} \end{aligned}$$

Theorem 1.5 (Theorem (4.5) in Patil (1993), and Theorem (4.5) in Patil-Singh (1990)) The set

$$\Omega = \{\xi_{i,j} \mid 1 \leq i \leq j \leq p-1\} \cup \{\theta\} \cup \{\varphi_i \mid i \in I\} \cup \{\psi_j \mid j \in J\}$$

forms a minimal generating set for the ideal P . Also, the set

$$\{\xi_{i,j} \mid 1 \leq i \leq j \leq p-1\} \cup \{\theta\} \cup \{\varphi_i \mid 0 \leq i \leq p-r\} \cup \{\psi_j \mid 0 \leq j \leq p-r'\}$$

forms a generating set for the ideal P .

Sengupta (2003) tried to prove that the set Ω from Theorem 1.5 forms a Groebner basis for P with respect to the grevlex monomial order using the grading $wt(x_i) = m_i$ with $x_0 < x_1 < \dots < x_n$. In this case $\prod_{i=0}^n x_i^{a_i} >_{\text{grevlex}} \prod_{i=0}^n x_i^{b_i}$ if in the ordered tuple $(a_1 - b_1, \dots, a_n - b_n)$ the left-most nonzero entry is negative. Sengupta's proof works for arithmetic sequences, but it is incomplete for the almost arithmetic sequences:

Remark 1.6 Assume $r' \geq r$, $\mu = 0$, and $W \neq \phi$. Then Patil generators are not a Groebner basis with respect to the grevlex monomial ordering with $x_0 < x_1 < \dots < x_n$ and with the grading $wt(x_i) = m_i$.

Proof. As $u - z = (q - q_z)p + (r - r_z)$ then $r' \geq r$ if and only if $r_z \geq r$. Assume $r' \geq r$, then $r_z - r + 1 > 0$ and also $\theta = x_n^v - x_0^\mu x_{p+r-r'} x_p^{q-q'-1}$. Assume also that $\mu = 0$ and $W \neq \phi$, then $I = [\max(r_z - r + 1, 0), p - r] = [r_z - r + 1, p - r]$. Under these assumptions the S-polynomial $S(\psi_k, \theta)$ can not be reduced to zero modulo Ω : for $0 \leq k < r_z - r + 1$ consider $S(\psi_k, \theta) = x_0^\mu S_1$ where $S_1 = x_0^{\lambda-1} x_k x_n^w - \underline{x_{r'+k} x_{p+r-r'} x_p^{q-1}}$, with the leading monomial underlined. We note that $LM(S_1)$ is a multiple of $LM(\xi_{r'+j, p+r-r'})$ only. Hence, the only possible way to reduce S_1 with respect to Ω is by using $\xi_{r'+j, p+r-r'}$. However, none of the terms of the binomial $S_1 + x_p^{q-1} \xi_{r'+j, p+r-r'} = x_{r+k} x_p^q - x_0^{\lambda-1} x_k x_n^w$ is a multiple of any of the leading terms of Patil generators. Therefore, it can not be reduced to 0 modulo Ω . ■

It remains to give a concrete example for which $r' \geq r$, $\mu = 0$, and $W \neq \phi$ so that Patil generators are not a Groebner basis:

Remark 1.7 Let $m_0 \geq 5$ be an odd integer. Let P be the defining ideal of the monomial curve that corresponds to the almost arithmetic sequence $m_0, m_0 + 1, m_0 - 1$. Then Patil generators for the ideal P are not a Groebner basis with respect to the grevlex monomial ordering with $x_0 < x_1 < \dots < x_n$ and with the grading $wt(x_i) = m_i$.

Proof. Observe: $p = 1, n = 2$, and $g_i = i(m_0 + 1)$ for all i .

Let v, μ , and z be as defined in Lemma 1.2. Then $v(m_0 - 1) = \mu m_0 + z(m_0 + 1)$ for some integers $\mu, z \geq 0$. This implies $\mu + z < v$. Note that

$$v(m_0 - 1) = \mu m_0 + z(m_0 + 1) = (\mu + z)(m_0 - 1) + \mu + 2z.$$

Thus $\mu + 2z = s(m_0 - 1)$ for some $s \geq 1$. Hence,

$$v > \mu + z \geq \frac{\mu}{2} + z = \frac{s}{2}(m_0 - 1) \geq \frac{m_0 - 1}{2}.$$

Thus,

$$v \geq \frac{m_0 + 1}{2}. \quad (1)$$

On the other hand, note that

$$\frac{m_0 + 1}{2}(m_0 - 1) = \frac{m_0 - 1}{2}(m_0 + 1) \in \Gamma'. \quad (2)$$

Therefore, by the minimality of v we must have

$$v \leq \frac{m_0 + 1}{2}. \quad (3)$$

By (1) and (3) it follows that $v = \frac{m_0 + 1}{2}$.

Let u, λ, w , and g_u be as defined in Lemma 1.2. Note

$$\frac{m_0 + 1}{2}(m_0 + 1) - m_0 = \frac{m_0 - 1}{2}(m_0 - 1) + m_0 \in \Gamma. \quad (4)$$

Therefore,

$$u \leq \frac{m_0 + 1}{2}. \quad (5)$$

Claim $w > 0$: if $w = 0$ then $g_u = \lambda m_0$, thus $u(m_0 + 1) = \lambda m_0$. But m_0 and $m_0 + 1$ are relatively prime, therefore, we must have $u = b m_0$ for some $b \geq 1$, a contradiction to (5). Thus $w > 0$.

Claim $\lambda < u$: by Lemma 1.2 we have $u(m_0 + 1) = \lambda m_0 + w(m_0 - 1)$. If $\lambda \geq u$ then $w(m_0 - 1) = u(m_0 + 1) - \lambda m_0 = u + (u - \lambda)m_0$, which implies $u \geq m_0 - 1$ as $w > 0$, a contradiction to (5). Thus Claim $\lambda < u$.

Now consider

$$\begin{aligned} w(m_0 - 1) &= u(m_0 + 1) - \lambda m_0, \\ &= (u - \lambda)(m_0 - 1) + 2u - \lambda. \end{aligned}$$

As $w(m_0 - 1) > 0$ and $u > \lambda$ we must have $2u - \lambda = c(m_0 - 1)$ for some $c \geq 1$. But if $u \leq \frac{m_0 - 1}{2}$ then $2u - \lambda \leq m_0 - 1 - \lambda$, a contradiction as $\lambda \geq 1$. Therefore,

$$u > \frac{m_0 - 1}{2}. \quad (6)$$

By (5) and (6) it follows that $u = \frac{m_0 + 1}{2}$.

Now by the uniqueness in Lemma 1.2 and as of (2) and (4) it follows that $\mu = 0$, $z = \frac{m_0 - 1}{2}$, $\lambda = 2$ and $w = \frac{m_0 - 1}{2}$.

Finally, note that $r = p = r' = 1$. Therefore, the parameters z, w, μ, p, r , and r' all satisfy the assumptions of the previous remark, hence done. ■

Example 1.8 Let $m_0 = 7, m_1 = 8$, and $m_2 = 6$. By the previous remark $v = 4$, $z = 3$, $\mu = 0$, $u = 4$, $\lambda = 2$, and $w = 3$. Therefore, as $p = 1$ then $q = 3, r = 1, q' = 0, r' = 1, q_z = 2$, and $r_z = 1$. Thus $I = [\max(r_z - r + 1, 0), p - r] = \emptyset$ and $J = [0, \min(z - 1, p - r')] = \{0\}$. By Patil (1993), or Theorem 1, a minimal set of generators for the ideal P is $\Omega = \{\psi_0, \theta\}$ where $\psi_0 = \underline{x_1 x_2} - x_0^2$ and $\theta = \underline{x_2^4} - x_1^3$. This is not a Groebner basis with respect the grevlex monomial order with $x_0 < x_1 < x_2$ and with the grading $\text{wt}(x_i) = m_i$: note neither term of $S(\psi_0, \theta) = x_1^4 - x_0^2 x_2^3$ is a multiple of $LT(\psi_0)$ or $LT(\theta)$.

2 Explicit Descriptions for I and J

Let $u, z, q, r, q' = q_{u-z}$, and $r' = r_{u-z}$ be as in Lemma 1.2 and Notation 1.3. Let $z = q_z p + r_z$ with $q_z \in \mathbb{Z}$ and $r_z \in [1, p]$. By Notation 1.1 it is clear that $q_z \leq q$ since $0 \leq z \leq u - 1$. As $u - z = (q - q_z)p + (r - r_z)$, it follows that $q' = q - q_z - \varepsilon$ and $r' = \varepsilon p + r - r_z$ where $\varepsilon = \begin{cases} 0, & \text{if } r > r_z; \\ 1, & \text{if } r \leq r_z. \end{cases}$

Therefore, $r' < r$ if and only if $r_z < r$. We note the following: if $r' < r$ then $r > r_z$, hence $r - r' = r_z$ and $q - q' = q_z$. Also, if $r' \geq r$ then $r \leq r_z$, hence $p + r - r' = r_z$ and $q - q' - 1 = q_z$. Therefore,

$$\theta = x_n^v - x_0^\mu x_{r_z} x_p^{q_z}.$$

Proposition 2.1 *Let $z > 0$ and let $z = q_z p + r_z$ with $q_z \in \mathbb{Z}$ and $r_z \in [1, p]$. Then*

$$\min\{z - 1, p - r'\} = \begin{cases} p - r', & \text{if } r \leq r_z; \\ p - r', & \text{if } r > r_z \text{ and } z > p; \\ z - 1, & \text{if } r > r_z \text{ and } z \leq p. \end{cases}$$

Moreover, $z \leq p$ if and only if $q_z = 0$.

Proof. First note that $p - r' = (1 - \varepsilon)p + r_z - r$. It is obvious that if $z > 0$ then $q_z \geq 0$. Consider three cases:

Case $r \leq r_z$: since $r \in [1, p]$ then $z - 1 = q_z p + r_z - 1 \geq r_z - 1 \geq r_z - r = p - r'$.

Case $r > r_z$ and $z > p$: this implies $q_z \geq 1$. Therefore, $z - 1 = q_z p + r_z - 1 \geq p + r_z - 1 \geq p + r_z - r = p - r'$.

Case $r > r_z$ and $z \leq p$: this implies $q_z = 0$. Therefore, $z - 1 = r_z - 1 \leq r_z - 1 + p - r < p + r_z - r = p - r'$. ■

Therefore, for $W \neq \phi$ we write J as following

$$J = \begin{cases} [0, p - r'], & \text{if } q_z > 0 \text{ or } \varepsilon > 0; \\ [0, r_z - 1], & \text{if } q_z = 0 \text{ and } \varepsilon = 0. \end{cases}$$

3 Groebner Basis

We will prove that the binomials listed here form a Groebner basis for the defining ideal P with respect to the grevlex monomial order with $x_0 < x_1 < \dots < x_n$ and with the grading $wt(x_i) = m_i$:

$$\begin{aligned} \varphi_i &= \underline{x_{r+i} x_p^q} - x_0^{\lambda-1} x_i x_n^w, & \text{for } 0 \leq i \leq p - r; \\ \psi_j &= \underline{x_{r'+j} x_p^{q'}} x_n^{v-w} - x_0^{\nu-1} x_j, & \text{for } j \in J \\ \theta &= \underline{x_n^v} - x_0^\mu x_{r_z} x_p^{q_z}, \\ \xi_{i,j} &= \begin{cases} \underline{x_i x_j} - x_0 x_{i+j}, & \text{if } i + j \leq p; \\ \underline{x_i x_j} - x_{i+j-p} x_p, & \text{if } i + j > p, \end{cases} & \text{for } 1 \leq i \leq j \leq p - 1, \end{aligned}$$

where the underlined monomials are the leading monomials.

Before we state the theorem that contains the main result of this article, let us recall the definition of the minimal Groebner basis.

Definition 3.1 A *minimal Groebner basis* for a polynomial ideal I is a Groebner basis G for I such that:

- (i) $LC(f) = 1$ for all $f \in G$, where $LC(f)$ is the leading coefficient of f .
- (ii) For all $f \in G$, $LM(f) \notin \langle LM\{G - \{f\}\} \rangle$, where $LM\{G - \{f\}\}$ is the set of leading monomials of all polynomials in the set $\{G - \{f\}\}$.

Now we state the main theorem of this article considering the binomials φ_i , ψ_j , θ , and $\xi_{i,j}$ as stated above.

Theorem 3.2 The set

$$\Phi = \{\varphi_i \mid 0 \leq i \leq p-r\} \cup \{\psi_j \mid j \in J\} \cup \{\theta\} \cup \{\xi_{i,j} \mid 1 \leq i \leq j \leq p-1\}$$

is a minimal Groebner basis for the ideal P with respect to the grevlex monomial order with $x_0 < x_1 < \dots < x_n$ and with the grading $wt(x_i) = m_i$.

The remainder of this article is to prove Theorem 3.2. The proof is a consequence of a general result that we state and prove in the main proposition of this article, Proposition 3.6. The following three lemmas are needed to prove the main proposition.

Lemma 3.3 Let $e_0, \dots, e_p, d_1 \in \mathbb{N}_0$ with $e_1 + \dots + e_p \geq 1$. The monomial $\alpha = x_0^{e_0} \dots x_p^{e_p} x_n^{d_1}$ can be reduced to $x_0^h x_s x_p^l x_n^d$ (modulo Φ) with respect to the grevlex monomial order such that $l \leq q$, $d < v$, $s \in [1, p]$, $h \geq e_0$, and with $\sum_{i=0}^p e_i m_i + d_1 m_n = h m_0 + m_s + l m_p + d m_n$.

Proof. Note that, using θ as many times as necessary, the monomial $\alpha = x_0^{e_0} \dots x_p^{e_p} x_n^{d_1}$ can be reduced to $x_0^{e'_0} \dots x_p^{e'_p} x_n^k$ with $k < v$ and $e'_1 + \dots + e'_p \geq e_1 + \dots + e_p$. This reduces the proof to the case $\alpha = x_0^{e_0} \dots x_p^{e_p} x_n^{d_1}$ with $d_1 < v$.

Using various $\xi_{i,j}$, α can be reduced to $\sigma = x_0^{h_1} x_{s_1} x_p^{l_1} x_n^{d_1}$ for some $h_1, l_1 \in \mathbb{N}_0$, and $s_1 \in [1, p]$ with $\sum_{i=0}^p e_i m_i = h_1 m_0 + m_{s_1} + l_1 m_p$. Observe that $g_{l_1 p + s_1} = l_1 m_p + m_{s_1}$.

If $l_1 \leq q$ then there is nothing to prove. If $l_1 > q$ apply the following algorithm:

Input $\sigma_1 = x_0^{h_1} x_{s_1} x_p^{l_1} x_n^{d_1}$.

Output $\sigma = x_0^h x_s x_p^l x_n^d$ with $l \leq q$, $d < v$, and $s \in [1, p]$ such that

$$hm_0 + m_s + lm_p + dm_n = h_1 m_0 + m_{s_1} + l_1 m_p + d_1 m_n.$$

Let $i = 0$.

REPEAT

Let $i = i + 1$.

Step 1: Write $l_i = a_i(q + 1) + b_i$ with $a_i \in \mathbb{N}_0$ and $b_i \in [0, q]$.

Reduce σ_1 with respect to φ_{p-r} a_i times:

$$\sigma' = x_0^{a_i(\lambda-1)+h_i} x_{s_i} x_{p-r}^{a_i} x_p^{b_i} x_n^{a_i w + d_i}.$$

Step 2: Write $a_i w + d_i = c_i v + d_{i+1}$ with $c_i \in \mathbb{N}_0$ and $d_{i+1} \in [0, v - 1]$ (note that $c_i \leq a_i$ because $w < v$ and $d_i < v$. Also, note that if $c_i = a_i$ then $d_{i+1} < d_i$).

Reduce σ' with respect to θ c_i times:

$$\sigma'' = x_0^{a_i(\lambda-1)+h_i+c_i\mu} x_{s_i} x_{r_z}^{c_i} x_{p-r}^{a_i} x_p^{c_i q_z + b_i} x_n^{d_{i+1}}.$$

Step 3: Reduce σ'' with respect to $\xi_{r_z, p-r}$ (or ξ_{p-r, r_z} if $p - r \leq r_z$. Recall that $\xi_{p-r, r_z} = x_0^{1-\varepsilon} x_{p-r'} x_p^\varepsilon$ and $p - r' = (1 - \varepsilon)p + r_z - r$) c_i times (recall $c_i \leq a_i$):

$$\begin{aligned} \sigma''' &= x_0^{a_i(\lambda-1)+h_i+c_i\mu} x_{s_i} \left(x_0^{1-\varepsilon} x_{p-r'} x_p^\varepsilon\right)^{c_i} x_{p-r}^{a_i-c_i} x_p^{c_i q_z + b_i} x_n^{d_{i+1}}, \\ &= x_0^{a_i(\lambda-1)+h_i+c_i\mu} x_{s_i} \left(x_0^{1-\varepsilon} x_p^\varepsilon\right)^{c_i} x_{p-r'}^{c_i} x_{p-r}^{a_i-c_i} x_p^{c_i q_z + b_i} x_n^{d_{i+1}}. \end{aligned}$$

Reduce σ''' with respect to various $\xi_{i,j}$ we get:

$$\sigma_{i+1} = x_0^{h_{i+1}} x_{s_{i+1}} x_p^{l_{i+1}} x_n^{d_{i+1}}.$$

with (by degree count) $l_{i+1} \leq c_i q_z + b_i + a_i + \varepsilon c_i$, and $h_{i+1} \geq a_i(\lambda - 1) + h_i + c_i \mu$, and some $s_{i+1} \in [1, p]$.

UNTIL $l_{i+1} \leq q$.

Note that $l_{i+1} \leq l_i$ since $c_i \leq a_i$ and since

$$l_{i+1} \leq c_i q_z + b_i + a_i + \varepsilon c_i \leq a_i(q - \varepsilon) + b_i + (1 + \varepsilon)a_i = a_i(q + 1) + b_i = l_i. \quad (7)$$

Now we prove that the sequence $l_1, l_2, \dots$ is a decreasing sequence which eventually goes below q :

Case $c_i < a_i$, $q_z = 0$ and $\varepsilon = 0$: then

$$l_{i+1} \leq c_i q_z + b_i + a_i + \varepsilon c_i = b_i + a_i < a_i(q+1) + b_i = l_i.$$

Case $c_i < a_i$, $q_z \neq 0$ or $\varepsilon \neq 0$: then (recall that $q > 0$ by Remark 1.4) then $l_{i+1} \leq l_i$ by (7).

Case $c_i = a_i$ for some $i > 0$: then we prove that there must be an integer $t > i$ such that $c_t < a_t$ and hence $l_{t+1} < l_t$: note that if $c_i = a_i$ then $d_i > d_{i+1}$. Hence $d_i, d_{i+1}, \dots$ is a strictly decreasing sequence of integers whenever $c_i = a_i, c_{i+1} = a_{i+1}, \dots$ since the d_i are non-negative hen necessarily there exist $t > i$ such that $c_t < a_t$. Therefore, $l_{t+1} < l_t$. ■

Lemma 3.4 Assume $W \neq \phi$. Let $\sigma = x_0^{h_1} x_{s_1} x_p^{l_1} x_n^{d_1}$ with $l_1 > q'$, $d_1 \geq v - w$, $h_1 \geq 0$, and $s_1 \in [1, p]$. Then σ can be reduced (modulo Φ) with respect to the grevlex monomial order to $x_0^h x_s x_p^l x_n^d$ with $l \leq q$, $d < v$, $h \geq h_1$, and $s \in [1, p]$ such that $h_1 m_0 + m_{s_1} + l_1 m_p + d_1 m_n = h m_0 + m_s + l m_p + d m_n$, with either $l \leq q'$ or $d < v - w$.

Proof. For the case $q_z = 0$ and $\varepsilon = 0$ we have $q' = q$ as $q' = q - q_z - \varepsilon$, therefore the proof is done by the previous lemma. Thus, we only need to work the proof for the case $q_z > 0$ or $\varepsilon > 0$. By the previous lemma the proof is reduced to $\sigma = x_0^{h_1} x_{s_1} x_p^{l_1} x_n^{d_1}$ with $l_1 \leq q$ and $d_1 < v$. Consider the following algorithm:

Input $\sigma_1 = x_0^{h_1} x_{s_1} x_p^{l_1} x_n^{d_1}$, $l_1 \leq q$, $d_1 < v$.

Output $\sigma = x_0^h x_s x_p^l x_n^d$ with $l \leq q'$ or $d < v - w$, and $s \in [1, p]$.

Set $i = 1$.

WHILE $l_i > q'$ and $d_i \geq v - w$ **REPEAT**:

Step 1: Write $d_i = a_i(v - w) + b_i$ with $a_i \in \mathbb{N}$ and $b_i \in [0, v - w - 1]$ (note $a_i > 0$ since $d_i \geq v - w$).

Write $l_i = a'_i(q' + 1) + b'_i$ with $a'_i \in \mathbb{N}$ and $b'_i \in [0, q']$ (note $a'_i > 0$ since $l_i > q'$).

Let $k_i = \min\{a_i, a'_i\}$. Note $k_i > 0$ since $a_i > 0$ and $a'_i > 0$.

Reduce σ_i with respect to $\psi_{p-r'}$ k_i times:

$$\sigma' = x_0^{h_i + k_i(v-1)} x_{s_i} x_{p-r'}^{k_i} x_p^{l_i - k_i(q'+1)} x_n^{d_i - k_i(v-w)}.$$

Step 2: Reduce σ' with respect to various $\xi_{a,b}$ we get:

$$\sigma_{i+1} = x_0^{h_{i+1}} x_{s_{i+1}} x_p^{l_{i+1}} x_n^{d_{i+1}}.$$

with some $l_{i+1} \leq l_i - k_i q' \leq l_i$ (since $k_i > 0$), $d_{i+1} = d_i - k_i(v - w) < d_i$, $h_{i+1} \geq h_i$, and some $s_{i+1} \in [1, p]$.

Let $i = i + 1$.

END LOOP

It is clear that the sequence $l_1, l_2, \dots$ is decreasing and the sequence $d_1, d_2, \dots$ is strictly decreasing. Hence the above algorithm must terminate. ■

Lemma 3.5 *Assume $W \neq \phi$ and $q' = 0$. Let $\sigma = x_0^{h_1} x_{s_1} x_n^{d_1}$ with $d_1 < v$, $h_1 \geq 0$, and $s_1 \in [1, p]$. Then σ can be reduced to $x_0^h x_s x_n^d$ with $(s, d) \in V - W$ and $h \geq h_1$.*

Proof. As $q' = 0$ then $\psi_j = \frac{x_{r'+j} x_n^{v-w}}{x_0^{\nu-1}} x_j$. Also, either $q_z > 0$ or $\varepsilon > 0$ since $q > 0$ and $q' = q - q_z - \varepsilon$, so that $J = [0, p - r']$. Let $a = \min\{t \geq 1 \mid s_1 - tr' < r'\}$ and $b = \min\{t \geq 1 \mid d_1 - t(v - w) < v - w\}$. Let $k = \min\{a, b\}$. Reduce σ with respect to $\psi_{s_1 - ir'}$ for $i = 1, \dots, k$ we get $x_0^{h+k(\nu-1)} x_{s_1 - kr'} x_n^{d_1 - k(v-w)}$, with either $s_1 - kr' < r'$ (hence $s_1 - kr' < u - z$) or $d_1 - k(v - w) < v - w$. Therefore, $(s_1 - kr', d_1 - k(v - w)) \in V - W$ as $s_1 \leq p < u$ and $d_1 - k(v - w) < v$ for $k \geq 1$ since $d_1 < v$. ■

Proposition 3.6 *Let $e_0, \dots, e_p, d_1 \in \mathbb{N}_0$ with $e_1 + \dots + e_p \geq 1$. The monomial $\alpha = x_0^{e_0} \dots x_p^{e_p} x_n^{d_1}$ can be reduced to $x_0^h x_s x_p^l x_n^d$ (modulo Φ) with respect to the grevlex monomial order, such that $\sum_{i=0}^p e_i m_i + d_1 m_n = h m_0 + m_s + l m_p + d m_n$ with $(lp + s, d) \in V - W$.*

Proof. We consider two cases $W = \phi$ and $W \neq \phi$ separately:

$W = \phi$: $V - W = V$. By Lemma 3.3 the proof is reduced to the case $\sigma = x_0^h x_s x_p^l x_n^d$, with $l \leq q$, $d < v$, and $s \in [1, p]$. If $lp + s < u$ then $(lp + s, d) \in V$ (since $d < v$), hence done. Otherwise, if $lp + s \geq u = qp + r$ then $l = q$ and $s \geq r$. Reduce σ using φ_{s-r} to $\sigma_2 = x_0^{h+\lambda-1} x_{s-r} x_n^{d+w}$. If $d + w < v$ then done since $(s - r, d + w) \in V$ (since $s - r < p < u$). Otherwise, if $d + w \geq v$ then $w > 0$, thus $z = 0$ as $W = \phi$, hence $q_z = -1$ and $r_z = p$, thus $\theta = x_n^v - x_0^\mu$. Reduce σ_2 using θ to $x_0^{h+\lambda-1+\mu} x_{s-r} x_n^{d+w-v}$. Observe $d + w - v < v$ since $d < v$ and $w < v$, hence $(s - r, d + w - v) \in V$ (since $s - r < p < u$), hence done.

$W \neq \phi$: by Lemma 3.4 the proof is reduced to the case $\sigma = x_0^h x_s x_p^l x_n^d$, with $l \leq q$, $d < v$, $s \in [1, p]$, and with either $l \leq q'$ or $d < v - w$. Consider the following cases:

- (1) $lp + s < u - z$: then $(lp + s, d) \in V - W$ (since $d < v$), hence done.
- (2) $lp + s \geq u - z$, consider three subcases:
 - (2a) $d < v - w$ and $lp + s < u$: then $(lp + s, d) \in V - W$, hence done.

(2b) $d < v - w$ and $lp + s \geq u$: as $u = qp + r$ and $l \leq q$ then $l = q$ and $s \geq r$. Reduce $\sigma = x_0^h x_s x_p^l x_n^d$ using φ_{s-r} to $\sigma' = x_0^{h+\lambda-1} x_{s-r} x_n^{d+w}$. Note $(s-r, d+w) \in V$ because $s-r < p < u$ and $d+w < v$. If $(s-r, d+w) \in V-W$ then done. Otherwise, if $(s-r, d+w) \notin V-W$ then necessarily $s-r \geq u-z = q'p + r'$, therefore, $q' = 0$ because $s-r < p$. Hence, done by Lemma 3.5.

(2c) $d \geq v - w$: then by application of Lemma 3.4, necessarily $l \leq q'$. Then $q'p + r' = u - z \leq lp + s \leq q'p + s$, so that $s \geq r'$ and $l = q'$. If we know that $s - r' \in J$, we could use $\psi_{s-r'}$ to reduce σ . According to that we consider two subcases:

(2c-1) $q_z = 0$ and $\varepsilon = 0$: this implies $q = q'$, therefore $q' > 0$. Under the assumptions, we have $l = q' = q$. Consider two cases:

Case $s \geq r$: reducing σ with respect to φ_{s-r} we get $\sigma' = x_0^{h+\lambda-1} x_{s-r} x_n^{d+w}$. As $d+w \geq v$ reduce σ' with respect to θ we get $\sigma'' = x_0^{h+\lambda+\mu-1} x_{s-r} x_{r_z} x_n^{d+w-v}$. Note $s-r+r_z < p$ as $\varepsilon = 0$ ($r > r_z$). Hence, using ξ_{s-r, r_z} (or $\xi_{r_z, s-r}$ if $r_z \leq s-r$) σ'' reduces to $x_0^{h+\lambda+\mu} x_{s-r+r_z} x_n^{d+w-v}$ with $(s-r+r_z, d+w-v) \in V-W$ since $s-r+r_z < p < u-z = q'p + r$ as $q' > 0$ and $d+w-v < v$. Hence, done.

Case $s < r$: in this case $0 \leq s-r' < r-r' = r_z$. Thus, $s-r' \in J$. Reduce σ with respect to $\psi_{s-r'}$ we get $x_0^{h+\nu-1} x_{s-r'} x_n^{d-(v-w)}$ with $s-r' < p < u-z = q'p + r$ as $q' > 0$. Hence, $(s-r', d-(v-w)) \in V-W$ as $d-(v-w) < v$. Hence, done.

(2c-2) $q_z > 0$ or $\varepsilon > 0$: hence $J = [0, p-r']$. If $q' = 0$ then done by Lemma 3.5. If $q' > 0$ then reduce σ using $\psi_{s-r'}$ to $x_0^{h+\nu-1} x_{s-r'} x_n^{d-(v-w)}$. Note $(s-r', d-(v-w)) \in V-W$ as $d-(v-w) < v$ and $s-r' < p < u-z = q'p + r'$ since $q' > 0$. Hence, done. ■

Proposition 3.7 Let $\alpha = x_0^{e_0} \cdots x_p^{e_p} x_n^h$ and $\beta = x_0^{d_0} \cdots x_p^{d_p} x_n^k$ with $e_0, \dots, e_p, h, d_0, \dots, d_p, k \in \mathbb{N}_0$, with $e_1 + \dots + e_p \geq 1$ and $d_1 + \dots + d_p \geq 1$ such that $\sum_{i=0}^p e_i m_i + h m_n = \sum_{i=0}^p d_i m_i + k m_n$. Then $\alpha - \beta$ can be reduced to zero modulo Φ .

Proof. By Proposition 3.6 α can be reduced to $x_0^{c'} x_{s'} x_p^{l'} x_n^{h+d'}$ with $c' \geq 0, (l'p + s', d') \in V-W$, and $s' \in [1, p]$. Furthermore, working with the same procedures as in the proof of Proposition 3.6 the monomial $x_0^{c'} x_{s'} x_p^{l'} x_n^{h+d'}$ can be reduced to $x_0^c x_s x_p^l x_n^d$ with $c \geq 0, (lp + s, d) \in V-W$, and $s \in [1, p]$. Note that we have $cm_0 + lm_p + m_s + dm_n = \sum_{i=0}^p e_i m_i + h m_n = \sum_{i=0}^p d_i m_i + k m_n$. Hence, the proof follows by Lemma 1.2 part (b). ■

Proof. (of the main theorem, Theorem 3.2) We apply Buchberger algorithm to show that all the S-polynomials, $S(f, g)$ with $f, g \in \Phi$, reduces to 0 with respect to Φ (modulo Φ). Note that, by Lemma 1.2, all the binomials in the set Φ are

homogeneous with respect to the grading $wt(x_i) = m_i$. Thus all the S-polynomials are homogeneous. Also, it is easy to see that each such S-polynomial has the form $\alpha - \beta$ as in Proposition 3.7. This shows that Φ is a Groebner basis for the ideal inP .

To complete the proof of Theorem 3.2 we need to prove the minimality of Φ . The proof is as follows: it is clear that $LM(\theta) \notin \langle LM(G - \{\theta\}) \rangle$. By Lemma 1.2 $w < v$ and by Remark 1.4 $q > 0$, hence it is clear that $LM(\varphi_i) \notin \langle LM(G - \{\varphi_i\}) \rangle$ and $LM(\xi_{i,j}) \notin \langle LM(G - \{\xi_{i,j}\}) \rangle$. If $q_z > 0$ or $\varepsilon > 0$ then $LM(\psi_j) \notin \langle LM(G - \{\psi_j\}) \rangle$ since $q' < q$ because $q' = q - q_z - \varepsilon$. If $q_z = 0$ and $\varepsilon = 0$ then $\{LM(\psi_j) \mid 0 \leq j \leq r_z - 1\} = \{x_j x_p^q x_n^{v-w} \mid r - r_z \leq j \leq r - 1\}$, hence $LM(\psi_j) \notin \langle LM(G - \{\psi_j\}) \rangle$. ■

Patil and Singh (1990) constructed a generating set (but not minimal) for the defining ideal P . We call the elements of this set “Patil-Singh generators”. This set of generators consists of the same binomials in the set of Patil generators but with adding a few more generators. The Patil-Singh generators are as follows: let q, r, q_z, r_z, q', r' , and ε and be as before,

$$\begin{aligned} \varphi_i &= \overline{x_{i+r} x_p^q} - x_0^{\lambda-1} x_i x_n^w, & \text{for } 0 \leq i \leq p-r; \\ \psi_j &= \overline{x_{r'+j} x_p^{q'} x_n^{v-w}} - x_0^{\nu-1} x_j, & \text{for } 0 \leq j \leq p-r'; \\ \theta &= \overline{x_n^v} - x_0^\mu x_{r_z} x_p^{q_z}, \\ \xi_{i,j} &= \begin{cases} \overline{x_i x_j} - x_0 x_{i+j}, & \text{if } i+j \leq p; \\ \overline{x_i x_j} - x_{i+j-p} x_p, & \text{if } i+j > p, \end{cases} & \text{for } 1 \leq i \leq j \leq p-1. \end{aligned}$$

The following theorem follows directly from Theorem 3.2.

Theorem 3.8 *The set $\{\varphi_i \mid 0 \leq i \leq p-r\} \cup \{\theta\} \cup \{\xi_{i,j} \mid 1 \leq i \leq j \leq p-1\} \cup \{\psi_j \mid 0 \leq j \leq p-r'\}$ is a Groebner basis for the ideal P with respect to the grevlex monomial order with $x_0 < x_1 < \dots < x_n$ with the grading $wt(x_i) = m_i$.*

Note that Theorem 3.8 gives a Groebner basis with an easier description but it gives up the minimality since if $q_z = 0$ and $\varepsilon = 0$, then $r_z < r_z + p - r = p - r'$ and $\{LM(\psi_j) = x_{j+r'} x_p^q x_n^{v-w} \mid r_z \leq j \leq p - r'\} = \{x_j x_p^q x_n^{v-w} \mid r \leq j \leq p\} = x_n^{v-w} \{LM(\varphi_i) = x_j x_p^q \mid 0 \leq j \leq p - r\}$.

Finally, we finish this article by noting that Patil-Singh generators do not form a Groebner basis in all cases if we consider the grevlex monomial order with the same grading as before but with $x_0 > x_1 > \dots > x_n$ (in this case $\prod_{i=0}^n x_i^{a_i} >_{grevlex} \prod_{i=0}^n x_i^{b_i}$ if in the ordered tuple $(a_1 - b_1, \dots, a_n - b_n)$ the right-most nonzero entry is negative). In the following we prove this and give an example.

Remark 3.9 Assume $r < r_z < p$ (hence $\varepsilon = 0$), $\lambda > 1$, and $w > 0$. Then Patil-Singh generators are not a Groebner basis with respect to the grevlex monomial ordering with $x_0 > x_1 > \dots > x_n$ and with the grading $wt(x_i) = m_i$.

Proof. First note the leading terms of the generators: $LT(\varphi_i) = x_{i+r}x_p^q$ if $w > 0$ and $LT(\varphi_i) = x_0^{\lambda-1}x_i$ if $w = 0$, $LT(\psi_j) = x_0^{\lambda+\mu-\varepsilon}x_j$, $LT(\theta) = x_0^\mu x_{r_z} x_p^{q_z}$, and $LT(\xi_{i,j}) = x_i x_j$. Assume $r < r_z < p$ (hence $\varepsilon = 0$), $\lambda > 1$, and $w > 0$. Consider $S(\xi_{1,r_z}, \theta) = x_1 x_n^v - x_0^{\mu+1} x_{r_z+1} x_p^{q_z}$. Under the above assumptions it is clear that none of the terms of $S(\xi_{1,r_z}, \theta)$ is a multiple of any of the leading terms of the Patil-Singh generators. ■

Example 3.10 Let $m_0 = 20, m_1 = 21, m_2 = 22, m_3 = 23, m_4 = 24$, and $m_5 = 29$. Hence $p = 4$. Let P be the kernel of the K -algebra homomorphism $\eta : K[x_0, \dots, x_5] \rightarrow K[t]$, defined by $\eta(x_i) = t^{m_i}$. Recall the parameters in Lemma 1.2. It is easy to check that $v = 3$, hence by the uniqueness condition we must have $\mu = 2$, $q_z = 1$, and $r_z = 3$, thus $z = 7$. For $1 \leq i \leq 3$ note that in order for $am_4 + m_i - m_0$ to be in Γ we must have $a \geq 2$. Note $g_{2p+1} = 2(24) + 21 = 2(20) + 29$. Therefore, we conclude that $q = 2$ and $r = 1$, thus $u = 9$. Hence, $\lambda = 2$, $w = 1$, $r' = 2$, and $q' = 1$. Therefore, Patil-Singh generators are as follows: $G = \{\varphi_i \mid 0 \leq i \leq 3\} \cup \{\psi_j \mid 0 \leq j \leq 2\} \cup \{\theta\} \cup \{\xi_{i,j} \mid 1 \leq i \leq j \leq 3\}$ where $\varphi_i = \underline{x_{i+1}x_4^2} - x_0 x_i x_5$, and $\psi_j = x_{j+2}x_5^2 - \underline{x_0^3 x_j}$, and $\theta = x_5^3 - \underline{x_0^2 x_3 x_4}$ and $\xi_{i,j}$ as defined before with $p = 4$. The set G is not Groebner basis with respect to the grevlex monomial order with $x_0 > x_1 > \dots > x_5$ and with the grading $wt(x_i) = m_i$: consider $S(\theta, \xi_{1,3}) = x_1 x_5^3 - x_0^3 x_4^2$. Note that neither term of $S(\theta, \xi_{1,3})$ is a multiple of any of the leading terms above.

Acknowledgement

The author thanks Prof. Swanson I. for the useful discussions and comments during the course of this work.

References

- [1] Cox, D., Little, J., O'Shea, D. (1996). *Ideals, Varieties and Algorithms*. New York: Springer-Verlag.
- [2] Greuel, G-M., Pfister, G. (2002). *A Singular Introduction to Commutative Algebra*. Berlin Heidelberg: Springer-Verlag.

- [3] Hübl, R., Swanson, I. (2002). Normal cones of monomial primes. *Mathematics of Computation* 72(241):459-475.
- [4] Patil, D. P. (1993). Minimal sets of generators for the relation ideals of certain monomial curves. *Manuscripta Math.* 80:239-248.
- [5] Patil, D. P., Singh, Balwant (1990). Generators for the derivation modules and the relation ideals of certain curves. *Manuscripta Math.* 68:327-335.
- [6] Sengupta, I. (2003). A Gröbner basis for certain affine monomial curves. *Comm. Algebra* 31(3):1113-1129.

Department of Mathematics and Statistics
Jordan University of Science and Technology
P O Box 3030, Irbid 22110, Jordan
Email address: iayyoub@just.edu.jo